

GROUP-SPECIFIC BACTERIAL DIVERSITY IN THE GUT OF FUNGUS-CULTIVATING TERMITES (*MACROTERMES*, *ODONTOTERMES* AND *MICROTERMES* SPECIES) FROM KENYA

H. M. Makonde^{1,2,3}, H. Peter-Klenk¹ and H. I. Boga^{2,4}

¹Leibniz-Institut DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Inhoffenstraße 7b, 38124 Braunschweig, Germany

²Jomo Kenyatta University of Agriculture and Technology, Nairobi, Kenya

³Technical University of Mombasa, Mombasa, Kenya

⁴Taita Taveta University College, Voi, Kenya

E-mail:- huxcly@yahoo.com

Abstract

The mutualistic association between termites and their gut symbionts has continued to attract the curiosity of researchers over time. The aim of this study was to characterize group-specific bacterial community structure and diversity in the gut of three fungus cultivating termites *Macrotermes michaelseni*, *Odontotermes* and *Microtermes* species using PCR-denaturing gradient gel electrophoresis (PCR-DGGE) analysis. Group-specific primers targeting members of the phyla *Bacteroidetes*, *Firmicutes* and *Planctomycetes* were used for PCR amplification of total DNA mixture extracted from termite guts. The PCR products were used as templates in a second PCR with nested bacteria DGGE-PCR primer pairs whose products were separated by DGGE. Representative DNA bands were excised from the gels, re-amplified, purified and sequenced. The sequences were blast analyzed and together with other reference sequences retrieved from the public GenBank were used to infer phylogenetic trees. All sets of sequences were deposited in the public GenBank. Results from the DGGE band patterns revealed a sharp contrast between the bacterial communities of *M. michaelseni*, *Odontotermes* and *Microtermes* species. This underlines the difference in group-specific bacterial diversity in the three termites. Phylogenetic analysis of the 16S rRNA gene sequences indicated that they were affiliated with the three phyla: *Bacteroidetes*, *Firmicutes* and *Planctomycetes*. Sequences (40%) affiliated with the phylum *Planctomycetes* were isolated and clustered with 'Termite *planctomycete* cluster', indicating that they are termite gut specific members. Sequences (89%) isolated were often affiliated with sequences obtained from other termites' guts, demonstrating that a majority of the gut bacteria are autochthonous having mutualistic relationship with their hosts. Notably, the isolated sequences had less than 96% sequence similarity with the closest cultivated strains, indicating that the majority of termite gut bacterial lineages are still uncultured. The results will help better understand the bacterial symbionts-termites mutualistic associations.

Key words: Bacterial diversity, termites, PCR-DGGE

1.0 Introduction

Termites play diverse roles in semi-arid and humid ecosystems and have an impact on the turnover of organic matter in tropical and subtropical regions. Termites rely on microbes in their guts for degradation of recalcitrant components of plant biomass and this has a major influence on soil structure and carbon mineralization (Brune and Friedrich, 2000; Brune and Ohkuma, 2011). The isolation and cultivation of several bacterial strains from termite guts has partially enabled their classification as decomposers of lignocellulose, uric acid and/or other aromatic compounds, as nitrogen-fixers, and/or as H₂/CO₂-acetogens (Breznak, 2000). Nonetheless, the majority of the microbial species are difficult or impossible to cultivate, thus limiting our knowledge on their role in the termite gut (Breznak, 2000). Therefore, alternative culture-independent methods can be used to increase our knowledge on the symbiont-host relationships.

Different approaches including 16S rRNA gene profiling have helped determining the microbial diversity in termites without cultivation (Hongoh *et al.*, 2003, 2005, 2006; Shinzato *et al.*, 2005, 2007; Fisher *et al.*, 2007; Long *et al.*, 2010; Mackenzie *et al.*, 2010; Mathew *et al.*, 2012; Köhler *et al.*, 2012). These culture-independent studies reported high bacterial diversity in the guts of termites including some termite-specific bacterial lineages (Hongoh *et al.*, 2003; Shinzato *et al.*, 2005). Recently, metagenomics studies have greatly expanded our knowledge of gut

symbiosis by providing key information on bacterial genes required for reductive acetogenesis, fermentation, lignocellulosic digestion, and nitrogen fixation within the host-symbiont association (Warnecke *et al.*, 2007; Tartar *et al.*, 2009; Mattéotti *et al.*, 2011; Liu *et al.*, 2011; Köhler *et al.*, 2012). However, there is little information on the evolutionary relationships between symbionts and host termites and the specificity of the bacterial lineages to their hosts. This has partly been attributed by the high abundance, cryptic and unpredictable foraging patterns (Ahmed *et al.*, 2011) coupled with the enormous diversity of their gut symbionts (Makonde *et al.*, 2013a).

The fungus cultivating termites comprise many of the economically important termite species (Ahmed *et al.*, 2011). Although comprehensive studies have been conducted on these termites (Hongoh *et al.*, 2006; Shinzato *et al.*, 2007; Long *et al.*, 2010; Mackenzie *et al.*, 2010; Mathew *et al.*, 2012; Zhu *et al.*, 2012, Makonde *et al.*, 2013a), comparative analysis of group specific bacterial lineages of the genera *Odontotermes*, *Macrotermes* and *Microtermes* has not been reported. This is important not only for detecting host specific bacterial lineages, but also for generating data that can be used to infer host-symbionts evolutionary relationships. This study, therefore, attempted to characterize group-specific bacterial community structure and diversity in the gut of three fungus cultivating termites *Macrotermes michaelseni*, *Odontotermes* and *Microtermes* species using PCR-denaturing gradient gel electrophoresis (PCR-DGGE) analysis. The results will contribute knowledge of the specificity and mutualistic relationship between gut bacterial symbionts and termites.

2.0 Material and Methods

2.1 Collection and Identification of Termites

Samples were collected in March, 2011 from Juja in Kiambu County, Kenya (latitude 1° 5' 54.68" N, longitude 37° 1' 1.10" W). Termite mounds (Designated as mound B colonized by *Macrotermes* and *Microtermes* species at the bottom and upper parts, respectively and mound C, approximately 2 km far apart was colonized by *Odontotermes* sp.) were excavated to a depth of 0.5 – 1.0 m. Termites (n = 200 workers and 50 soldiers) were sampled into sterile plastic boxes. Worker-caste termites were used in the experiments due to their foraging behavior during establishment and renewal of the fungus gardens. The identity of the termites was confirmed by sequencing the mitochondrial cytochrome oxidase II gene in DNA extracted from the heads of soldiers (Austin *et al.*, 2004; Makonde *et al.*, 2013b) and comparing it to the sequences of previously identified specimens.

2.2 DNA Extraction and PCR Amplification

The exterior surfaces of the termites were washed with 70% ethanol and then rinsed with sterile distilled water. The guts were aseptically removed with forceps (Schmitt-Wagner *et al.*, 2003). A total of 26 guts (approximately 144 mg) from *Macrotermes mchaelseni* [JQ247993] and *Odontotermes* sp. [JQ247986] and 74 guts (approximately 143 mg) from *Microtermes* sp. [JQ247990] were put separately into three sterile micro tubes containing 0.2 ml of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0). They were then homogenized using a sterile glass rod and used for total DNA extraction as described elsewhere (Makonde *et al.*, 2013b) using the UltraClean® Mega soil DNA isolation kit (MO BIO Laboratories, Inc.) according to the manufacturer's protocol.

2.3 Group-specific PCR-DGGE Analysis

The primers used were those described in other studies (Muyzer *et al.*, 1993, 1998; Mühling *et al.*, 2008). Group-specific primers (Table 1) were used to PCR amplify total DNA from guts of *Macrotermes michaelseni*, *Microtermes* and *Odontotermes* species. For each PCR, 1 µl (25 ng/ µl) of the template was mixed with TaKaRa Ex Taq™ HS (5 units/ µl) according to the manufacturer's protocol. The PCR conditions were as described by Mühling *et al.*, 2008 and the PCR product size was checked on 1% agarose gel stained with ethidium bromide.

PCR products were used as templates in a second PCR with nested bacteria DGGE-PCR primer pairs. PCR conditions were as described by Mühling *et al.*, (2008) but with distinct annealing temperatures (AT) as indicated in Table 1. PCR products were separated by DGGE in an 8% (w/v) polyacrylamide gel with urea and formamide as denaturants using an Ingeny phorU system (Ingeny International BV, Goes, The Netherlands). Linear denaturing gradients between 40% and 60% (Mühling *et al.*, 2008) or 35% to 80% were used. Electrophoresis was performed in 1x TAE buffer at 60° C at an initial voltage of 200 V for 5 min, followed by a constant voltage of 100 V for 16 hours.

Polyacrylamide gels were stained with SYBRGold (MoBiTec, Göttingen, Germany) for 1 hour. DNA bands of interest were excised from the gels with a sterile scalpel, and the DNA was eluted overnight at 4° C in 25 µl of 10 mM Tris-HCl buffer (pH 8.0).

2.4 PCR re-amplification and Sequencing

Eluates (3 µl) were used for re-amplification as described previously using corresponding primers without a GC clamp. PCR products were directly purified and sequenced at Helmholtz Centre for Infection Research (HZI), Braunschweig, Germany. Trace files were manually edited and assembled using Invitrogen vector NTI 11.5 software. Bands that resulted in poor sequences were cloned into pGEM-T[®] Easy vector system II (Promega) according to the manufacturer's recommendations and transfected through heat shock into *E. coli* JM109 high efficiency competent cells (Promega). Selection of transformants and extraction of plasmid DNA followed described protocols (Ausubel, 1995). Representative clones were selected and then sequenced as described above. All sets of sequences were deposited in GenBank under the accession numbers JX421956 to JX421963 (DGGE *Bacteroidetes* sequences), JX421964 to JX421986 (DGGE *Firmicutes* sequences) and JX421987 to JX422007 (DGGE *Planctomycetes* sequences).

2.5 Phylogenetic Analysis

The sequences were quality checked for chimeric structures using the Mallard program (Ashelford *et al.*, 2006). A search for similar sequences using BLASTN (Altschul *et al.*, 1990) was performed, and sequence alignment was performed using the CLUSTAL Omega program (<http://www.clustal.org>) against the nearest neighbours. A neighbor-joining tree of the aligned sequences was constructed (Saitou and Nei, 1987) using MEGA V5.10 (Tamura, 2011). Evolutionary distances were computed using the Maximum Composite Likelihood method (Tamura, 2004). Bootstrapping (Felsenstein, 1985) was conducted with 1000 replicates to obtain statistical support values. All sites, including gaps in the sequence alignment, were excluded pairwise in the phylogenetic analysis. The taxonomic assignment was confirmed at an 90% confidence level using the naïve Bayesian rRNA classifier on the RDP website (Cole *et al.*, 2005).

3.0 Results

3.1 Nested PCR-DGGE Analysis

Comparative analysis of the DGGE profiles from the gut contents of three termites (*Macrotermes michaelseni*, *Odontotermes* and *Microtermes* spp.) displayed visual differences in the DGGE band patterns (Figure 1). *Bacteroidetes* DGGE profiles had poor resolution and only three bands for *Odontotermes* sp. could be resolved. Likewise five bands for *Microtermes* sp. were analyzed (Figure 1A). Members from the phylum *Bacteroidetes* could not be resolved by DGGE analysis in *M. michaelseni* thus no clear bands were observed. The *Firmicutes* DGGE profile for *Microtermes* sp. displayed nine bands but only seven bands could be analyzed (Fig. 1B). Likewise, the DGGE profile for *Odontotermes* sp. had seven bands with different intensities, which were analyzed while nine bands for *M. michaelseni* were analyzed. Notably, band numbers 9, 10, 15, 16 and 17 were shared (Fig. 1B). The *Planctomycetes* DGGE profiles showed seven bands for *Odontotermes* sp., but displayed eight bands for *Microtermes* sp. and six bands for *M. michaelseni*. However, band numbers 24, 25, 28, 29 and 36 were shared by either two or all termites (Figure 1C, Table 2).

3.2 Phylogenetic Affiliation of Sequences

The 16S rDNA sequence analysis of the bands excised from DGGE gels revealed that the bands were affiliated with the phyla *Bacteroidetes*, *Firmicutes* and *Planctomycetes*. Phylogenetic analysis of the sequences grouped the respective bacterial phyla together with corresponding sequences previously isolated from termite guts (Figure 2A, B & C). Most of the newly isolated *Bacteroidetes* sequences were closely affiliated (97-99% sequence similarity) with other sequences obtained from gut of termites. The sequence DGGE8B8 [JX421763] was phylogenetically closely related to bacterial clones isolated from cockroach and *Pachnoda ephipiata* [FJ374177] (with 96-97% sequence identity) while sequences DGGE8B5-6 [JX421959-61] had less than 96% sequence similarity with clone sequences from sediments cockroach and human feces (Figure 1A; Table 2). This scenario was also observed for

the newly isolated *Firmicutes* and *Planctomycetes* sequences where majority of them were related to other sequences, which were often obtained from other termites (Table 2).

In the Phylogenetic tree of the *Firmicutes* (Figure 2B), two main clusters were formed. One cluster had three isolated sequences (DGGEFB9, DGGEF10 and DGGEF22) that were affiliated with *Lactococcus lactis* and other clones isolated from vegetable, termite gut and mouse (sequence identity of <97% with a bootstrap value of 99%). The other seven isolated sequences (DGGEFB12, DGGEFB14, DGGEFB15, DGGEFB16, DGGEFB17, DGGEFB21 and DGGEFB23) formed the second cluster (with a bootstrap value of 100%) and were often affiliated (sequence identity of 96-98%) with clone sequences from *Nasutitermes takasagoensis* and *Cubitermes orthognathus*. For the phylogenetic tree of the phylum *Planctomycetes*, only a single main cluster was formed with all newly isolated sequences affiliated (95-99% sequence similarity) with *Planctomycete* clones from different termite guts (*Microcerotermes*, *Cubitermes* spp., *Coptotermes formosanus* and *Reticulitermes* spp.) as indicated in Figure 2C and Table 2. Some of the sequences (highlighted in green) were shared between the termites. This indicates the presence of termite gut specific bacterial lineages previously observed in the phylum *Firmicutes*. The blast comparison and phylogenetic analysis also indicated that the co-migrating bands were 100% identical (Figure 2A, B & C; Table 2).

4.0 Discussion

The phylogenetic analysis of the group-specific bacterial 16S rRNA genes from the three termites revealed bacterial communities that are still uncultured. The affiliation of the isolated DGGE sequences with others previously obtained from termite guts underscores the existence of termite specific bacterial lineages (Hongoh *et al.*, 2006; Shinzato *et al.*, 2007; Warnecke *et al.*, 2007; Makonde *et al.*, 2013a). The co-migration of some bands indicated that some of the bacterial lineages are shared in these termites. This scenario, however, can also emanate from either random acquisition of microorganisms from the environment (Curtis *et al.*, 2004) or variation in the hosts' diets (Tanaka *et al.*, 2006). Members from the phylum *Bacteroidetes* could not be resolved by DGGE in *M. michaelseni*. This could be due to poor separation of the bands as a result of the denaturation gradient (35-80%) used. The presence of unique DGGE bands in each termite may demonstrate the presence of termite genera specific bacterial lineages.

Previously, members of the phylum *Bacteroidetes* were detected as the dominant phylogenetic groups using 16S rRNA clone based methods (Hongoh *et al.*, 2006; Shinzato *et al.*, 2007; Mackenzie *et al.*, 2010; Zhu *et al.*, 2012, Makonde *et al.*, 2013a). However, the low resolution in the DGGE analysis limited the detection of majority members, which demonstrates the pitfall of the DGGE method in *Bacteroidetes* profiling. Notably, members of *Bacteroidetes* have been shown to play important roles in the termites; for instance, members of the genera *Prevotella* (*Prevotella ruminicola* 23 and *Prevotella intermedia* 17) can ferment both xylan and cellulose through carbohydrate-active enzymes such as xylanase, carboxymethylcellulase and endoglucanase (<http://www.cazy.org>). Kodama *et al.* (2012) reported a *Bacteroidetes* isolate, *Dysgonomonas oryzae* from a microbial fuel cell, which implicates it in cellulose degradation. Genome studies have revealed *Bacteroides* involvement in biodegradation of polysaccharides and fermentation of sugars (Xu *et al.*, 2003; Sonnenburg *et al.*, 2010) by contributing glycosyl hydrolases for their hosts' digestion (Liu *et al.*, 2011). Some members of *Bacteroidetes* also benefit their host by excluding potential pathogens from colonizing the gut (Wexler, 2007).

The DGGE profile for the phylum *Firmicutes* indicated bands with different intensities, demonstrating difference in relative abundances of the respective bacterial members. Moreover, it shows the DGGE method can better resolve members of the *Firmicutes* (Fig. 2B). The unique DGGE bands (DGGEFB12-14 [421970-72], DGGEFB21-23 [421984-86]) detected in each termite gut emphasize specific bacterial-hosts associations while the presence of co-migration bands (DGGEFB9 [421964-66], DGGEFB10 [421967-68], DGGEFB15 [421973-75], DGGEFB16 [421976-78], DGGEFB17 [421979-80]) underlines the termite gut specific bacterial lineages previously reported (Hongoh *et al.*, 2006; Shinzato *et al.*, 2007; Warnecke *et al.*, 2007; Makonde *et al.*, 2013a). Most of the newly isolated sequences were closely related to other clone sequences from termites. The termite gut specific bacteria may have important role for the survival of the termites. Moreover cultivated members of *Firmicutes* such as *Acetivibrio*

cellulosolvens, *Acetivibrio multivorans*, *Acetivibrio ethanolgignens* and *Acetivibrio cellulolyticus* (Tanaka *et al.*, 1991; Xu *et al.*, 2003) have been reported to ferment cellulosic substrates.

Notably, using the group-specific PCR-DGGE fingerprinting method, more 16S rRNA gene sequences affiliated with the phylum *Planctomycetes* were captured compared to previous clone-based analysis (Mackenzie *et al.*, 2010; Zhu *et al.*, 2012, Makonde *et al.*, 2013a) which comprehensively described the dominant phyla. All of the *Planctomycete* sequences clustered with those from other termites and belong to a large clade, the 'Termite *Planctomycete* cluster' (Köhler *et al.*, 2008), indicating that they are termite gut specific members. Again, this demonstrates that the low abundance of *Planctomycetes* previously detected in the gut of other fungus-cultivating termites (Shinzato *et al.*, 2005; Hongoh *et al.*, 2006; Mackenzie *et al.*, 2010; Long *et al.*, 2010; Zhu *et al.*, 2012) could be as a result of PCR bias (von Wintzingerode *et al.*, 1997). This, however, needs to be substantiated with more samples using general bacterial primers and group-specific primers.

5.0 Conclusion

The DGGE analysis confirmed the presence of common gut bacterial species in the investigated termites (*M. michaelsoni*, *Odontotermes* and *Microtermes* spp.) that could be important for the termites' life styles. The presence of unique DGGE banding patterns for each termite may indicate termite genus-specific bacterial lineages. Though the method used in this study cannot help infer physiological roles for the uncultured bacteria in the termites, the findings provide more knowledge on specific bacterial community structure in the guts of fungus-cultivating termites and contribute to understanding gut bacterial diversity and their associations with termites.

Acknowledgement

This work was supported by a PhD scholarship from the Deutscher Akademischer Austauschdienst (DAAD) and a grant (NCST/5/003/PG/224) from the Kenya National Council for Science and Technology (NCST). Kenya Wildlife Services (KWS) and National Environmental Management Act (NEMA) provided permits for sample collection in Kenya.

References

- Ahmed, B. M., Nkunika, P. O. Y., Sileshi, W. G., French, J. R. J., Nyeko, P. and Jain, S. (2011). Potential Impact of Climate Change on Termite Distribution in Africa. *British Journal of Environment and Climate Change*, **1**, pp 172-189.
- Altschul, S. F., Gish, W. and Miller, W, et al. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, **215**, pp 403–410.
- Ashelford, K. E., Chuzhanova, N. A., Fry, J. C., Jones, A. J. and Weightman, A. J. (2006). New screening software shows that most recent large 16S rRNA gene clone libraries contain chimeras. *Applied Environmental Microbiology*, **72**, pp 5734-5741.
- Austin, J. W., Szalanski, A. L. and Cabrera, B. J. (2004). Phylogenetic analysis of the subterranean termite family *Rhinotermitidae* (Isoptera) by using the mitochondrial cytochrome oxidase II gene. *Annual Entomology Society of America*, **97**, pp 548-555.
- Ausubel, F. M. (1995). Current protocols in molecular biology. John Wiley and Sons, New York.
- Breznak, J. A. (2000). Ecology of prokaryotic microbes in the guts of wood- and litter-feeding termites. In: Abe T., Bignell D.E., Higashi M. (eds) *Termites: Evolution, Sociality, Symbiosis, Ecology*, Kluwer Academic Publishers, Dordrecht, pp 209-231.
- Brune, A. and Friedrich, M. (2000). Microecology of the termite gut: structure and function on a microscale. *Current Opinion in Microbiology*, **3**, pp 263-269.
- Brune A. and Ohkuma M. (2011). Role of the termite gut microbiota in symbiotic digestion. In: Bignell D.E., Roisin Y., Lo, N. (eds) *Biology of termites: a modern synthesis*, Springer, Dordrecht, pp 439-475.
- Cole, J. R., Chai, B. and Farris, R. J., et al. (2005). The Ribosomal Database Project (RDP-II): Sequences and tools for high-throughput rRNA analysis. *Nucleic Acids Response*, **33**, pp D294–D296.
- Curtis, T. P. and Sloan, W. T. (2004). Prokaryotic diversity and its limits: microbial community structure in nature and implications for microbial ecology. *Current Opinion in Microbiology*, **7**, pp 221-226.
- Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, **39**, pp 783-791.
- Fisher, M., Miller, D., Brewster, C., Husseneder, C. and Dickerman, A. (2007). Diversity of gut bacteria of *Reticulitermes flavipes* as examined by 16S rRNA gene sequencing and amplified rDNA restriction analysis. *Current Microbiology*, **55**, pp 254-259.
- Hongoh, Y., Deevong, P., Inoue, T., Moriya, S., Trakulnaleamsai, S., Ohkuma, M., Vongkaluang, C., Noparatnaraporn, N. and Kudo, T. (2005). Intra- and Interspecific Comparisons of Bacterial Diversity and Community Structure Support Coevolution of Gut Microbiota and Termite Host. *Applied Environmental Microbiology*, **71**, pp 6590-6599.
- Hongoh, Y., Ekpornprasit, L., Inoue, T., Moriya, S., Trakulnaleamsai, S., Ohkuma, M., Noparatnaraporn, N. and Kudo, T. (2006). Intracolony variation of bacterial gut microbiota among castes and ages in the fungus-growing termite *Macrotermes gilvus*. *Molecular Ecology*, **15**, pp 505-516.
- Hongoh, Y., Ohkuma, M. and Kudo, T. (2003). Molecular analysis of bacterial microbiota in the gut of the termite *Reticulitermes speratus* (Isoptera; Rhinotermitidae). *FEMS Microbiol Ecology*, **44**, pp 231-242.

Kodama, Y., Shimoyama, T. and Watanabe, K. (2012). *Dysgonomonas oryzaevi* sp. nov., isolated from a microbial fuel cell. *International Journal of Systematic and Evolution Microbiology*. doi:10.1099/ijs.0.039040-0.

Köhler, T., Dietrich, C., Scheffrahn, R. H. and Brune, A. (2012). High-resolution analysis of gut environment and bacterial microbiota reveals functional compartmentation in the gut of wood-feeding higher termites (*Nasutitermes* spp.). *Applied Environmental Microbiology*, **78**, pp 4691-4701.

Lee, S.H., Malone, C. and Kemp, P.F. (1993). Use of 16S rRNA targeted fluorescent probes to increase signal strength and measure cellular RNA from natural planktonic bacteria. *Mar Ecological Progress Series*. **101**, pp 193-201.

Liu, N., Yan, X., Zhang, M., Xie, L., Wang, Q., Huang, Y., Zhou, X., Wang, S. and Zhou, Z. (2011). Microbiome of fungus-growing termites: a new reservoir for lignocellulase genes. *Applied Environmental Microbiology*, **77**, pp 48-56.

Long, Y. H., Xie L., Liu, N., Yan, X., Li, M. H., Fan M. Z. and Wang, Q. (2010). Comparison of gut-associated and nest-associated microbial communities of a fungus-growing termite (*Odontotermes yunnanensis*). *Journal of Insect Science*, **17**, pp 265-276.

Mackenzie, L. M., Muigai, A. T., Osir, E. O., Lwande, W., Keller, M., Toledo, G. and Boga, H. I. (2007). Bacterial diversity in intestinal tract of fungus-cultivating termite *Macrotermes michaelseni* (Sjöstedt). *African Journal of Biotechnology*, **6**, pp 658-667.

Makonde, H. M., Boga, H. I., Osiemo, Z., Mwirichia, R., Stielow, B., Göker, M. and Klenk, H. P. (2013b). Diversity of *Termitomyces* associated with fungus-farming termites assessed by cultural and culture-independent methods. *PloS ONE*. 8: e56464, doi:10.1371/journal.pone.0056464.

Makonde, H. M., Boga, H. I., Osiemo, Z., Mackenzie, L. M., Göker, M. and Klenk, H. P. (2013a). 16S-rRNA-based analysis of bacterial diversity in the gut of fungus-cultivating termites (*Microtermes* and *Odontotermes* species). *Antonie van Leeuwenhoek*, **104**, pp 869-883.

Mathew, G. M., Ju, Y.-M., Lai C.-Y., Mathew D.C., Huang C.C. (2012). Microbial community analysis in the termite gut and fungus comb of *Odontotermes formosanus*: the implication of *Bacillus* as mutualists. *FEMS Microbiology Ecology*, **79**, pp 504-517.

Mattéotti C., Thonart P., Francis F., Haubruge E., Destain J., Brasseur C., Bauwens J., De Pauw E., Portetelle D. and Vandenberg M. (2011). New glucosidase activities identified by functional screening of a genomic DNA library from the gut microbiota of the termite *Reticulitermes santonensis*. *Microbiological Response*, **166**, pp 629-642.

Matthies, C., Gößner, A., Acker, G., Schramm, A. and Drake, H. L. (2004). *Lactovum miscens* gen. nov., sp. nov., an aerotolerant, psychrotolerant, mixed-fermentative anaerobe from acidic forest soil. *Research in Microbiology*, **155**, pp 847-854.


Mühling, M., Allen, W. J., Murrell, J. C. and Joint, I. (2008). Improved group-specific PCR primers for denaturing gradient gel electrophoresis analysis of the genetic diversity of complex microbial communities. *ISME J*, **2**, pp 379-392.


Muyzer, G., Brinkhoff, T., Nübel, U., Santegoeds, C., Schäfer, H. and Waver, C. (1998). Denaturing gradient gel electrophoresis (DGGE) in microbial ecology. In: Akkermans A.D.L., van Elsas J.D., de Bruijn F.J. (ed). *Molecular Microbial Ecology Manual*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 1-27.

- Muyzer, G., de Wall, E. C. and Uitterlinden A.G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied Environmental Microbiology*, **59**, pp 3233-3241.
- Saitou, N. and Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, **4**, pp 406-425.
- Schmitt-Wagner, D., Friedrich, M. W., Wagner, B. and Brune, A. (2003). Phylogenetic diversity, abundance, and axial distribution of bacteria in the intestinal tracts of two soil-feeding termites (*Cubitermes* spp). *Applied Environmental Microbiology*, **69**, pp 6007-6017.
- Shinzato, N., Muramatsu, M., Matsui, T. and Watanabe, Y. (2005). Molecular phylogenetic diversity of the bacterial community in the gut of the termite *Coptotermes formosanus*. *Biosciences Biotechnology and Biochemistry*, **69**, pp 1145-1155.
- Shinzato, N., Muramatsu, M., Matsui, T. and Watanabe, Y. (2007). Phylogenetic analysis of the gut bacterial microflora of the fungus-growing termite *Odontotermes formosanus*. *Biosciences Biotechnology and Biochemistry*, **71**, pp 906-915.
- Sonnenburg, E. D., Zheng, H., Joglekar, P., Higginbottom, S. K., Firbank, S. J., Bolam, D. N. and Sonnenburg J. L. (2010). Specificity of polysaccharide use in intestinal *Bacteroides* species determines diet-induced microbiota alterations. *Cell*, **141**, pp 1241–1252.
- Tamura, K., Nei, M. and Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbour-joining method. *Proceedings of National Academy of Science, USA*, **101**, pp 11030–11035.
- Tamura, K., Peterson, D., Peterson, N., Stecher G., Nei, M. and Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology Evolution*, **28**, pp 2731-2739.
- Tartar, A., Wheeler, M. M., Zhou, X., Coy, M. R., Boucias, D. G. and Scharf, M. E. (2009). Parallel metatranscriptome analyses of host and symbiont gene expression in the gut of the termite *Reticulitermes flavipes*. *Biotechnology Biofuels*, **2**, pp 25.
- Von Wintzingerode, F., Göbel, U. B. and Stackebrandt, E. (1997). Determination of microbial diversity in environmental samples: pitfalls of PCR-based rRNA analysis. *FEMS Microbiol Review*, **21**, pp 213-229.
- Warnecke, F., Luginbuhl P., Ivanova, N., Ghassemian, M., Richardson, T. H., Stege, J. T., Cayouette, M., McHardy, A. C. *et al.*, (2007). Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite. *Nature*. **450**, pp 560-565.
- Wexler, H. M. (2007). Bacteroides: the good, the bad, and the nitty-gritty. *Clinical Microbiology Review*. **20**, pp 593-621.
- Xu, J., Bjursell, M. K., Himrod, J., Deng, S., Carmichael, L. K., Chiang, H. C., Hooper, L. V. and Gordon, J. I. (2003). A genomic view of the human *Bacteroides thetaiotaomicron* symbiosis. *Science*, **299**, pp 2074-2076.
- Zhu, Y.Y., Li, J., Liu, H., Yang, H., Xin, S., Zhao, F., Zhang, X., Tian, Y. and Lu, X. (2012). Phylogenetic analysis of the gut bacterial microflora of the fungus-growing termite *Macrotermes barneyi*. *African Journal of Microbiology Research*, **6**, pp 2071-2078.


Figure 1

A. *Bacteroidetes*






B. Firmicutes



C. Planctomycetes

Figure 1: Nested PCR-DGGE band patterns for the phylum A. Bacteroidetes, B. Firmicutes and C. Planctomycetes for termites. Lanes: 'O' denotes *Odontotermes* sp., 'Ma' denotes *M. michaelsoni* and 'Mi' denotes *Microtermes* sp. Numbers and arrows indicate DNA bands that were excised and sequenced. Co-migrating bands are indicated by one number

Figure 2






Figure 2: Evolutionary relationships between partial 16S rRNA gene sequences and selected taxa in the phyla A. Bacteroidetes, B. Firmicutes and C. Planctomycetes. The DGGE sequences highlighted in green were shared by either two or the three termites. *Methanoculleus thermophilus* (accession number, AB065297) was used to root the trees

Table 1: Primers used for group-specific bacterial amplification. Primer sequences are described below

Target Group	Primers used for group-specific PCR _a	AT PCR (° C)	Primers used for re-PCR for DGGE	AT nested PCR (° C)	Denaturing gradient used for DGGE (%)
Firmicutes	Firm350f/814r	58	518f-GC/785r	57	40-60
Planctomycetes	Plancto352f/920r	68	518f-GC/907r	60	40-60
Bacteroidetes	CFB555f/968r	61	CFB555f-GC/907r	58	35-80

Key:

Firm350f: 5'-GGC AGC AGT RGG GAA TCT TC-3' (Mühling *et al.*, 2008),

Firm814r: 5'-ACA CYT AGY ACT CAT CGT TT-3' (Mühling *et al.*, 2008),

Plancto352f: 5'-GGC TGC AGT CGA GRA TCT-3' (Mühling *et al.*, 2008),

Plancto920r: 5'-TGT GTG AGC CCC CGT CAA-3' (Mühling *et al.*, 2008),

CFB555f: 5'-CCG GAW TYA TTG GGT TTA AAG GG-3' (Mühling *et al.*, 2008),

CBF968r: 5'-GGT AAG GTT CCT CGC GTA-3' (Mühling *et al.*, 2008),

907r: 5'-CCG TCA ATT CMT TTG AGT TT-3' (Muyzer *et al.*, 1993),

518f: 5'-CCA GCA GCC GCG GTA AT-3' (Muyzer *et al.*, 1993),

785r: 5'-CTA CCA GGT ATC TAA TCC-3' (Lee *et al.*, 1993).

GC clamp: CGC CCG CCG CGC GCG GCG GGC GGG GCG GGG GCA CGG GGG G (Muyzer *et al.*, 1998),

'AT' denotes annealing temperature.

Table 2: List of DGGE bands for termite guts 16S rRNA Gene sequences and their closest affiliations

Phylum	Band No: ^a	Accession number			Taxonomical Affiliation/ isolation source	Accession no.	% ID
		O	Ma	MI			
<i>Bacteroidetes</i>	1	-	-	JX421956	Clone Rs-E83 (<i>R. speratus</i>)	AB088937	99
	2	-	-	JX421957	Clone M1PL1-34 (<i>Microcerotermes</i> sp. 1)	AB191987	98
	3	-	-	JX421958	Clone RPK-74 (<i>Reticulitermes</i> sp.)	AB192268	96
	4	-	-	JX421959	Clone RII-AN131 (Sediment)	JQ580510	95
	5	-	-	JX421960	Clone SL42 (Cockroach)	JN680585	94
	6	JX421961	-	-	Bacterium NLAE-zl-H474 (Human feces)	JX006685	93
	7	JX421962	-	-	Clone SL41 (Cockroach)	JN680583	96
	8	JX421963	-	-	Bacterium clone PeHg27 (<i>P. ehippiata</i>)	FJ374177	97
<i>Firmicutes</i>	9	JX421964	JX421966	JX421965	Firmicutes clone Cf4-79 (<i>C. formosanus</i>)	GQ502559	95
	10	-	JX421968	JX421967	Bacterium clone F_NIH2_P08 (Mouse)	JF838107	94
	11	-	-	-	-	-	-
	12	JX421970	-	-	<i>Lactococcus lactis</i> (Vegetable)	AB571485	96

	13	-	JX4219	-	Bacterium clone COB P3-18 (<i>C. orthognathus</i>)	AY160826	96
			71				
	14	JX421972	-	-	Bacterium clone mbII-b2 (soil)	AB481151	98
	15	JX421975	JX4219	JX421974	Bacterium clone COB P3-18 (<i>C. orthognathus</i>)	AY160826	97
			73				
	16	JX421977	JX4219	JX421978	Bacterium clone COB P3-18 (<i>C. orthognathus</i>)	AY160826	96
			76				
	17	JX421979	-	JX421980	Bacterium clone COB P3-18 (<i>C. orthognathus</i>)	AY160826	98
	20	-	JX4219	-	Bacterium clone ECH_aai40c10 (<i>T. aculeatus</i>)	EU774894	96
			83				
	21	JX421984	-	-	Bacterium clone COB P3-18 (<i>C. orthognathus</i>)	AY160826	96
	22	-	-	JX421985	<i>Lactococcus lactis</i> (Vegetable)	AB571485	96
	23	-	-	JX421986	Bacterium clone COB P3-18 (<i>C. orthognathus</i>)	AY160826	97
Planctomyces	24	JX421987	-	JX421988	Planctomycete clone M1NP1-4 (<i>Microcerotermes</i> sp. 1)	AB192132	96
	25	JX421991	JX4219	JX421990	Planctomycete clone BCf2-25 (<i>C. formosanus</i>)	AB062813	97
			92				
	26	-	-	JX421993	Planctomycete clone Cf5-01 (<i>C. formosanus</i>)	GQ502587	99
	27	-	-	JX421994	Planctomycete clone BCf2-25 (<i>C. formosanus</i>)	AB062813	97
	28	JX421996	-	JX421995	Planctomycetales clone Rc152	JQ617850	97

					(<i>Reticulitermes</i> sp.)		
29	JX421998	JX4219	JX421997		Planctomycetales clone Rc152	JQ617850	97
		99			(<i>Reticulitermes</i> sp.)		
30	JX422000	-	-		Planctomycete clone Csp1104	AM77418	97
					(<i>Cubitermes</i> sp.)	7	
32	JX422002	-	-		Planctomycetales clone Rs-E96 (<i>R.</i>	AB089118	95
					<i>speratus</i>)		
33	JX422003	-	-		Planctomycete clone BCf2-25 (<i>C.</i>	AB062813	97
					<i>formosanus</i>)		
34	-	-	JX422004		Planctomycete clone RsW01-054 (<i>R.</i>	AB198483	95
					<i>speratus</i>)		
36	-	JX4220	JX422007		Planctomycetales clone Rs-E96 (<i>R.</i>	AB089118	97
		06			<i>speratus</i>)		

Key: 'DGGEBB' denotes denaturing gradient gel electrophoresis Bacteroidetes band, 'DGGEFB' denotes denaturing gradient gel electrophoresis Firmicutes band, and 'DGGEPB' denotes denaturing gradient gel electrophoresis Planctomycetes band. Number in bold indicates bands that were shared. '-' denotes absence of band, 'O' denotes *Odontotermes* sp., 'Ma' denote *Macrotermes michaelsoni* and 'Mi' denotes *Microtermes* sp.. ^aNumber in bold indicates bands that were shared. '-' denotes absence of band sequence.